

Presenting dcpos and dcpo algebras

Achim Jung*

M. Andrew Moshier[†]

Steven Vickers[‡]

March 17, 2008

Abstract

Dcpo can be presented by a preorder of generators and inequational relations expressed as covers. Algebraic operations on the generators (possibly with their results being ideals of generators) can be extended to the dcpo presented, provided the covers are “stable” for the operations. The resulting dcpo algebra has a natural universal characterization and satisfies all the inequational laws satisfied by the generating algebra.

Applications include known “coverage theorems” from locale theory.

1 Introduction

The importance of algebraic theories in semantics was first emphasized by Hennessy and Plotkin in [HP79] and more recently — in the context of “computational effects” — by Plotkin and Power, [PP04]. Mathematically, it necessitates the ability to construct free algebras over semantic spaces. For continuous domains, a general theory was developed by Abramsky and the first author in [AJ94, Section 6] making use of the concept of an “abstract basis.” There it was also stated that free algebras exist for the more general category **DCPO**, however, the construction was not concrete but reference was made to the *Freyd Adjoint Functor Theorem*. The result presented in this paper amends this situation and allows us to give a fairly concrete description of free dcpo algebras.

The key ingredient of our approach is a construction which produces a dcpo algebra from a preordered algebra subject to a certain set of “covers,” each of which forces a directed supremum to cover an element. Our first main theorem states that in the construction process one can ignore the operations of the given preordered algebra P and complete P simply with respect to the set C of covers; the operations can automatically be lifted to the resulting dcpo where they will be Scott-continuous. As our second main theorem we have the result that the dcpo algebra so constructed satisfies all inequations that hold in the original algebra P .

The proof techniques we employ are greatly inspired by those found in the paper [JV91] by Johnstone and the third author, where they were used to show that preframes can be presented by generators and relations.

The technique of presenting algebras by generators and relations is well known from Universal Algebra and works with magical ease, but a recurring difficulty is that it does not readily yield concrete information about the algebra presented.

*School of Computer Science, The University of Birmingham, Birmingham, United Kingdom; A.Jung@cs.bham.ac.uk

[†]Department of Mathematics & Computer Science, Chapman University, Orange, CA, USA; moshier@chapman.edu

[‡]School of Computer Science, The University of Birmingham, Birmingham, United Kingdom; s.j.vickers@cs.bham.ac.uk

One way to obtain partial information is through the *coverage theorems* of locale theory. We can illustrate this with the *Double Coverage Theorem* of [VT04, Theorem 7]. This is a very convenient illustration since we shall follow its techniques fairly closely, though it was by no means the first such coverage theorem. (We give a more extensive survey in Section 6, with the Double Coverage Theorem as Theorem 6.3.)

Suppose a frame is presented with a set of generators L that, as it happens, is a distributive lattice whose finite meets and joins are to be preserved in the frame. Suppose also R is a set of relations that are \vee -stable and \wedge -stable in a certain sense. Then

$$\mathbf{Fr}\langle L \text{ (qua DL)} \mid R \rangle \cong \mathbf{dcpo}\langle L \text{ (qua poset)} \mid R \rangle$$

Here “qua” means that the indicated structure is to be preserved. For example, “qua DL” (“DL” standing for distributive lattice) indicates that the injection

$$\eta: L \rightarrow \mathbf{Fr}\langle L \text{ (qua DL)} \mid R \rangle$$

preserves finite meets and joins. Similarly, “qua poset” indicates that the other injection of generators is monotone. In addition, the isomorphism commutes with those two injections.

Presenting frames by generators and relations is a central technique in locale theory. Logically, it amounts to locales *classifying* propositional geometric theories, in the same way as toposes can classify predicate geometric theories. The frame presentation gives us very sharp information about how to define frame homomorphisms out of the frame presented. (Locally, it tells us about locale maps into the corresponding locale, in other words the generalized points of the locale.) However, it does not immediately tell us the concrete structure of the frame. The Double Coverage Theorem tells us that if the presentation is sufficiently nice, then we know at least enough about the concrete structure to be able to define dcpo homomorphisms out of the frame. This is thus more extensive knowledge than we got immediately from the frame presentation.

The “niceness” of the presentation may look like a restriction, but in fact every presentation can be worked into the nice form by finitary means, by freely generating a distributive lattice from the generators.

Technically, the proof of the Double Coverage Theorem works in the following steps. First, show that the right-hand side really does present a dcpo; next, use its universal property to define meet and join operators; then, show that these are meet and join for the dcpo order, and are distributive. This shows that the right-hand side is in fact a frame. Next, to show the universal property specified by the frame presentation, it must be shown that certain dcpo homomorphisms are in fact frame homomorphisms.

A range of such coverage theorems are known, and can be proved by similar means. The aim of the present paper is to give a general account of the infinitary, dcpo aspects of this, showing how operations on generators and inequational laws holding for them can be extended to dcpos presented by generators and relations.

2 DCPO presentations

We first show that dcpo presentations by generators and relations do indeed present: in other words, there is a dcpo with the universal property specified in the presentation. This was essentially proved in [VT04, Section 2.1], using the techniques (originally used for preframes) of [JV91]. Our development here generalizes theirs in two ways. First, we assume that the generators form a preorder rather than a poset. This is inessential, since the preorder can be posetified. Second, [VT04] used equational relations of the form, “directed join = directed join.” We shall instead use inequational relations, of the form, “element \sqsubseteq directed join.” This is equivalent to the equational relations in the semilattice contexts of [VT04], but not more generally.

Definition 2.1 A dcpo presentation consists of

- a set P of generators;
- a preorder \sqsubseteq on P ;
- a subset C of $P \times \mathcal{P}(P)$, whose elements are called covers and written $a \triangleleft U$, subject to the requirement that U is directed with respect to \sqsubseteq . (Without the directedness requirement, we shall refer to C as a cover set.)

The goal is to show that every dcpo presentation gives rise to a dcpo with the appropriate universal property.

Definition 2.2 An order preserving map $f: (P; \sqsubseteq, C) \rightarrow D$ from a dcpo presentation to a dcpo D preserves covers if for all $a \triangleleft U$ in C it is true that $f(a) \sqsubseteq \bigsqcup_{x \in U} f(x)$.

Definition 2.3 A dcpo \bar{P} is freely generated by the dcpo presentation $(P; \sqsubseteq, C)$ if there is a map $\eta: P \rightarrow \bar{P}$ that preserves covers, and every map f from P to a dcpo D that preserves covers factors through η via a unique Scott-continuous map $\bar{f}: \bar{P} \rightarrow D$:

$$\begin{array}{ccc}
 \bar{P} & \xrightarrow{\bar{f}} & D \\
 \eta \uparrow & \nearrow f & \\
 P & &
 \end{array}$$

Definition 2.4 Let $(P; \sqsubseteq, C)$ be a preorder with cover set. A C -ideal I is a subset of P which is downward closed and closed under all covers, to wit, $U \sqsubseteq I$ implies $a \in I$ for all $a \triangleleft U$ in C .

If S is any subset of P then $\langle S \rangle$ denotes the smallest C -ideal containing S . The set of all C -ideals is denoted by $C\text{-Idl}(P)$.

This is the same definition as in [Joh82, Section II-2.11], however, our presenting preorder is not assumed to be a meet semilattice. Trivially, $C\text{-Idl}(P)$ is a complete lattice as $S \mapsto \langle S \rangle$ is a closure operator on the powerset of P . Specifically, $\bigsqcup_{k \in K} I_k = \langle \bigcup_{k \in K} I_k \rangle$ for any set $\{I_k \mid k \in K\}$ of C -ideals.

Now, $C\text{-Idl}(P)$ is *not* the free dcpo generated by a dcpo presentation but it plays a crucial role in our construction. Indeed, we shall be particularly interested in $C\text{-Idl}(P)$ as a sup-lattice (or complete join semilattice). As objects, sup-lattices and complete lattices are the same, but sup-lattice homomorphisms are only required to preserve all joins.

Proposition 2.5 Let (P, \sqsubseteq) be a preorder and C a cover set on it. Then $C\text{-Idl}(P)$ is

$$\text{SupLat}\langle P \text{ (qua preorder)} \mid a \sqsubseteq \bigsqcup U \text{ whenever } a \triangleleft U \rangle.$$

Proof. (From [AV93]) If Q is a sup-lattice and $f: P \rightarrow Q$ a monotone function that preserves covers, then define $\bar{f}: C\text{-Idl}(P) \rightarrow Q$ by

$$\bar{f}(I) = \bigsqcup_{p \in I} f(p).$$

This is clearly monotone and satisfies $\bar{f} \circ \eta = f$, so only the preservation of suprema needs to be shown. By monotonicity we have

$$\bar{f}\left(\bigsqcup_{k \in K} I_k\right) = \bigsqcup_{p \in \bigsqcup I_k} f(p) \supseteq \bigsqcup_{k \in K} \bigsqcup_{p \in I_k} f(p) =: x$$

For the other inequality note that $f^{-1}(\downarrow x)$ is downward closed by the monotonicity of f , closed under covers because these are assumed to be preserved by f , and a superset of all I_k . Hence $\bigsqcup_{k \in K} I_k = \langle \bigcup_{k \in K} I_k \rangle \subseteq f^{-1}(\downarrow x)$ and $\bar{f}(\bigsqcup_{k \in K} I_k) \sqsubseteq x$. \blacksquare

We shall also need the following lemma.

Lemma 2.6 *Let D be a dcpo. Then the sup-lattice $\Sigma(D)$ of Scott closed subsets of D is the free sup-lattice over D qua dcpo.*

Proof. A proof was sketched in [VT04] but we can also apply the preceding proposition by considering the dcpo presentation (D, \sqsubseteq, C) where C contains all covers $a \triangleleft U$ where $a \sqsubseteq \bigsqcup^\uparrow U$ in D . Then the set of C -ideals coincides with the set of Scott-closed subsets, and any monotone function $f: D \rightarrow E$ which preserves covers is in fact Scott-continuous. \blacksquare

Note that for a constructive treatment we must define ‘‘Scott closed’’ to mean down-closed and closed under directed joins, not the complement of a Scott open.

Note also that $\eta: D \rightarrow C\text{-Idl}(D) = \Sigma(D)$ takes each $a \in D$ to the corresponding principal ideal $\downarrow a$, and therefore is an order embedding. In the following we denote it by \downarrow , rather than η .

To return to the task of determining the dcpo generated by a given dcpo presentation, we let \bar{P} be the smallest sub-dcpo of $C\text{-Idl}(P)$ containing all $\langle p \rangle$ for $p \in P$, and define $\eta: P \rightarrow \bar{P}$ as $p \mapsto \langle p \rangle$. This is order-preserving because C -ideals are lower sets. For the preservation of covers let $a \triangleleft U$. Then $U \subseteq \bigcup_{x \in U}^\uparrow \langle x \rangle \subseteq \langle \bigcup_{x \in U}^\uparrow \langle x \rangle \rangle = \bigsqcup_{x \in U}^\uparrow \langle x \rangle$ and this contains a (and hence $\langle a \rangle$) because C -ideals are closed under covers.

Theorem 2.7 *\bar{P} together with the map $\eta: P \rightarrow \bar{P}$ is*

$$\text{dcpo}\langle P \text{ (qua preorder)} \mid a \sqsubseteq \bigsqcup U \text{ whenever } a \triangleleft U \rangle.$$

Proof. Let D be a dcpo and $f: P \rightarrow D$ a monotone function that preserves covers. The composite function $\downarrow \circ f: P \rightarrow \Sigma(D)$ also preserves covers and hence factors via a sup-lattice homomorphism $f': C\text{-Idl}(P) \rightarrow \Sigma(D)$. In other words, we have the following commuting diagram

$$\begin{array}{ccccc} P & \xrightarrow{\eta} & \bar{P} & \hookrightarrow & C\text{-Idl}(P) \\ & \searrow f & & & \downarrow f' \\ & & D & \hookrightarrow & \Sigma(D) \end{array}$$

and the idea is to pull back f' along \downarrow . For this note that the image of \downarrow is a sub-dcpo of $\Sigma(D)$ and since f' preserves suprema, $X_f := f'^{-1}(\downarrow(D))$ is a sub-dcpo of $C\text{-Idl}(P)$. Because the diagram commutes, $\eta(P)$ is a subset of X_f and then the same is true for \bar{P} . It follows that f' can be restricted to \bar{P} giving us a Scott-continuous map f'' from \bar{P} to $\downarrow(D) \subseteq \Sigma(D)$. Since $\downarrow(D)$ is order-isomorphic to D , $\bar{f} := \downarrow^{-1} \circ f''$ is a Scott-continuous map from \bar{P} to D with $f = \bar{f} \circ \eta$.

This is unique because if we had two such maps then their equalizer would be a sub-dcpo of \bar{P} that includes P and hence equal to \bar{P} . \blacksquare

It is worthwhile to point out that this result generalizes the usual ideal completion of a pre-order, which is obtained by letting the set of covers be empty. Similarly to the ideal completion it is true in the general case that although the empty set is always a C -ideal, it is never a member of \bar{P} .

We conclude this section with the following result which will turn out to be the crucial ingredient in our study of dcpo algebra presentations.

Proposition 2.8 *Let (P_i, \sqsubseteq, C_i) , $i = 1, \dots, n$, be a finite family of dcpo presentations. Then $\prod_i \bar{P}_i$ is the smallest sub-dcpo of $\prod_i C\text{-Idl}(P_i)$ containing $\prod_i \eta(P_i)$.*

Proof. Certainly $\prod_i \overline{P}_i$ is a sub-dcpo of $\prod_i C\text{-Idl}(P_i)$ that includes $\prod_i \eta(P_i)$. Now suppose D is any such sub-dcpo. Suppose for some i ($0 \leq i \leq n$) we have

$$\eta(P_1) \times \cdots \times \eta(P_i) \times \overline{P}_{i+1} \times \cdots \times \overline{P}_n \subseteq D.$$

This is certainly true for $i = n$. Let $S = \{a \in \overline{P}_i \mid \eta(P_1) \times \cdots \times \eta(P_{i-1}) \times \{a\} \times \overline{P}_{i+1} \times \cdots \times \overline{P}_n \subseteq D\}$. S is a sub-dcpo of \overline{P}_i that includes $\eta(P_i)$, and hence is the whole of \overline{P}_i . We deduce that $\eta(P_1) \times \cdots \times \eta(P_{i-1}) \times \overline{P}_i \times \cdots \times \overline{P}_n \subseteq D$, and it follows by induction on $n - i$ that $\prod_i \overline{P}_i \subseteq D$. \blacksquare

Since the transition from P to \overline{P} (with respect to a set C of covers) is the main operation in this paper, we give three alternative descriptions.

The d -topology Consider the d -topology on a dcpo having sub-dcpo's as its closed sets.¹ Then obviously \overline{P} is just the d -closure of the image $\eta(P)$. The two noteworthy properties of the d -topology are that it is Hausdorff on any dcpo ($\downarrow x$ is always open and closed), and that a map between dcpo's is Scott-continuous if and only if it is d -continuous and order-preserving.

The μ -topology Recall Keye Martin's μ -topology of a dcpo, generated by the subbasic open sets $\downarrow x \cap O$ with O a Scott-open subset, [Mar00, Chapter 3]. Alternatively, μ is the join of the lower Alexandrov and the Scott topologies.

Proposition 2.9 *The smallest sub-dcpo \overline{P} of a subset P of a dcpo D is contained in the μ -closure of P . Furthermore, for every element $x \in \overline{P}$ and Scott-open set O containing x there exists $x' \in P$ with $x' \in O$ and $x' \sqsubseteq x$.*

Proof. For the supremum of a directed set A to belong to a subbasic μ -open set $\downarrow x \cap O$, all elements of A must be below x . Because O is Scott-open, some element of A must meet it and this element will then be in $\downarrow x \cap O$. This shows that $D \setminus (\downarrow x \cap O)$ is a sub-dcpo of D , and hence that \overline{P} belongs to the μ -closure of P . \blacksquare

The two need not be the same, even when the ambient dcpo D is continuous:

Example 2.10 *Let D be the powerset of \mathbb{N} . For the elements of P choose the set of A_i , $i \in \mathbb{N}$ where $A_i = \mathbb{N} \setminus \{i\}$. These are pairwise incomparable, so the dcpo closure of P is P itself. On the other hand, every μ -neighbourhood of \mathbb{N} , the largest element of $\mathcal{P}\mathbb{N}$, contains a subset $\uparrow F$ with F a finite set. Clearly, $\uparrow F$ contains almost all A_i , so \mathbb{N} is in the μ -closure of P .*

Iterated ideal completion For every preorder P we have the ideal completion $\text{Idl}(P)$ consisting of *directed* lower sets of P . It is the free dcpo over P . Given a dcpo presentation $(P; \overline{\sqsubseteq}, C)$ we have the order-preserving map $\eta: P \rightarrow C\text{-Idl}(P)$ which therefore factors through the ideal completion $\text{Idl}(P)$: $\eta = \llbracket \uparrow \circ \downarrow \rrbracket$. Let's call the image P_1 . It is not necessarily a sub-dcpo of $C\text{-Idl}(P)$ but the process can be repeated: the inclusion of P_1 into $C\text{-Idl}(P)$ extends to a continuous map from $\text{Idl}(P_1)$ to $C\text{-Idl}(P)$. The image of this we call P_2 . And so on. At a limit ordinal λ we set $P_\lambda := \bigcup_{\alpha < \lambda} P_\alpha$. Eventually this sequence stabilizes at \overline{P} .

Each of these descriptions is useful in its own way but we must leave open the general problem of finding an *intrinsic* characterisation of those subsets of P that are elements of \overline{P} .

3 Operations

For $(P; \overline{\sqsubseteq}, C)$ a dcpo presentation, we now show how a monotone operation on P can be lifted to a Scott-continuous operation on the presented dcpo \overline{P} . As one may suspect, this will require a compatibility condition between the operation and the set of covers.

¹This appears to have been considered first by Oswald Wyler in [Wyl81].

So let $\omega: P^n \rightarrow P$ be an n -ary monotone operation on the preorder P . We say that C is *stable* for ω (or simply ω -*stable*) if for every $1 \leq i \leq n$, $a_i \triangleleft U$ in C , and $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in P$, the cover $\omega(a_1, \dots, a_n) \triangleleft \{\omega(a_1, \dots, a_{i-1}, x, a_{i+1}, \dots, a_n) \mid x \in U\}$ is also in C . All our results require stability and so from now on this is always assumed even if it is not explicitly stated.

The operation can be lifted to subsets in the obvious, pointwise fashion: For $A_i \subseteq P$ ($1 \leq i \leq n$) set

$$\omega(A_1, \dots, A_n) := \{\omega(a_1, \dots, a_n) \mid a_i \in A_i\}.$$

Lifting it to C -ideals requires the application of the associated closure operator: if $I_i \in C\text{-Idl}(P)$ ($1 \leq i \leq n$) set

$$\bar{\omega}(I_1, \dots, I_n) := \langle \{\omega(a_1, \dots, a_n) \mid a_i \in I_i\} \rangle.$$

Proposition 3.1 *For any $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in P$ and $T \subseteq P$ the following holds*

$$\omega(a_1, \dots, a_{i-1}, \langle T \rangle, a_{i+1}, \dots, a_n) \subseteq \langle \omega(a_1, \dots, a_{i-1}, T, a_{i+1}, \dots, a_n) \rangle.$$

Proof. Let us write S for the right-hand side above, and let

$$T' = \{x \in P \mid \omega(a_1, \dots, a_{i-1}, x, a_{i+1}, \dots, a_n) \in S\}.$$

By definition of S , we have $T \subseteq T'$. We show that T' is a C -ideal. Suppose $a \triangleleft U$ is a cover in C and $U \subseteq T'$. By stability, $\omega(a_1, \dots, a_{i-1}, a, a_{i+1}, \dots, a_n) \triangleleft \omega(a_1, \dots, a_{i-1}, U, a_{i+1}, \dots, a_n)$, and since the right-hand side belongs to the C -ideal S , so does the left-hand side. It follows that $a \in T'$ and therefore $\langle T \rangle \subseteq T'$. ■

Proposition 3.2 *As a function from $C\text{-Idl}(P)^n$ to $C\text{-Idl}(P)$, $\bar{\omega}$ preserves all joins in each argument.*

Proof. Clearly, $\bar{\omega}$ is monotone. Fixing some i , we must show

$$\bar{\omega}(I_1, \dots, \bigsqcup_{\lambda} J_{\lambda}, \dots, I_n) \subseteq \bigsqcup_{\lambda} \bar{\omega}(I_1, \dots, J_{\lambda}, \dots, I_n).$$

Let us write K for the right-hand side. Then we must show

$$\omega(I_1, \dots, \bigsqcup_{\lambda} J_{\lambda}, \dots, I_n) \subseteq K:$$

in other words, if, for each $j \neq i$, we have $a_j \in I_j$, then

$$\omega(a_1, \dots, \bigsqcup_{\lambda} J_{\lambda}, \dots, a_n) \subseteq K.$$

Putting $T = \bigcup_{\lambda} J_{\lambda}$ in Proposition 3.1, we have

$$\begin{aligned} \omega(a_1, \dots, \bigsqcup_{\lambda} J_{\lambda}, \dots, a_n) &\subseteq \langle \omega(a_1, \dots, \bigcup_{\lambda} J_{\lambda}, \dots, a_n) \rangle \\ &= \langle \bigcup_{\lambda} \omega(a_1, \dots, J_{\lambda}, \dots, a_n) \rangle \end{aligned}$$

Since K is a C -ideal and trivially for each λ ,

$$\omega(a_1, \dots, J_{\lambda}, \dots, a_n) \subseteq \omega(I_1, \dots, J_{\lambda}, \dots, I_n) \subseteq K.$$

we have

$$\langle \bigcup_{\lambda} \omega(a_1, \dots, J_{\lambda}, \dots, a_n) \rangle \subseteq K$$

which completes the argument. ■

As a corollary, $\bar{\omega}$ preserves all *directed* joins in each argument, and hence is jointly Scott continuous, and hence Scott continuous.

Lemma 3.3 *If $A_i \subseteq P$ ($1 \leq i \leq n$) then*

$$\bar{\omega}(\langle A_1 \rangle, \dots, \langle A_n \rangle) = \langle \omega(A_1, \dots, A_n) \rangle.$$

Proof. That the right-hand side is included in the left-hand side is just monotonicity. For the other inclusion we apply Proposition 3.1 n times to get the chain of inequalities

$$\omega(\langle A_1 \rangle, \langle A_2 \rangle, \dots, \langle A_n \rangle) \subseteq \langle \omega(A_1, \langle A_2 \rangle, \dots, \langle A_n \rangle) \rangle \subseteq \dots \subseteq \langle \omega(A_1, \dots, A_n) \rangle$$

Applying the C -ideal closure operator to the leftmost and rightmost term gives the result. |

Corollary 3.4 *The map $\eta: P \rightarrow C\text{-Idl}(P)$ preserves ω , in other words, if $a_1, \dots, a_n \in P$ then*

$$\bar{\omega}(\eta(a_1), \dots, \eta(a_n)) = \eta(\omega(a_1, \dots, a_n)).$$

Proposition 3.5 $\bar{\omega}$ maps \bar{P}^n to \bar{P} .

Proof. We must show that $\bar{P}^n \subseteq \bar{\omega}^{-1}(\bar{P})$. Since $\bar{\omega}$ is Scott continuous, $\bar{\omega}^{-1}(\bar{P})$ is a sub-dcpo of $C\text{-Idl}(P)^n$, and by Corollary 3.4 it includes $(\eta(P))^n$. Hence by Proposition 2.8 it includes \bar{P}^n . |

Theorem 3.6 *Let D be a dcpo and let $\omega_D: D^n \rightarrow D$ be a continuous map. Let further $f: P \rightarrow D$ be a monotone and cover preserving homomorphism. Then the extension $\bar{f}: \bar{P} \rightarrow D$ (defined in the proof of Theorem 2.7) is also a homomorphism.*

Proof. Consider the set

$$\{(I_1, \dots, I_n) \in \bar{P}^n \mid \omega_D(\bar{f}(I_1), \dots, \bar{f}(I_n)) = \bar{f}(\bar{\omega}(I_1, \dots, I_n))\}.$$

This is a sub-dcpo of \bar{P}^n because it is the equalizer of the two Scott-continuous functions $\omega_D \circ \bar{f}^n$ and $\bar{f} \circ \bar{\omega}$. It includes the image of P^n under η^n since

$$\begin{aligned} \omega_D(\bar{f}(\eta(a_1)), \dots, \bar{f}(\eta(a_n))) &= \omega_D(f(a_1), \dots, f(a_n)) \\ (f \text{ is a homomorphism}) &= f(\omega(a_1, \dots, a_n)) \\ &= \bar{f}(\eta(\omega(a_1, \dots, a_n))) \\ (\text{Corollary 3.4}) &= \bar{f}(\bar{\omega}(\eta(a_1), \dots, \eta(a_n))) \end{aligned}$$

Hence it equals \bar{P}^n . |

From a topological point of view, we can say the following about the algebra \bar{P} .

The d -topology Since a function between dcpos is Scott-continuous if and only if it is monotone and d -continuous, it then follows from Proposition 3.2 that $C\text{-Idl}(P)$ is a topological algebra with respect to the d -topology. Since $\eta(P)$ is a subalgebra of $C\text{-Idl}(P)$ it now follows for completely general reasons that its d -closure is a subalgebra as well. As noted at the end of Section 2, the d -closure is exactly \bar{P} .

The μ -topology We note that the operation $\bar{\omega}$ on $C\text{-Idl}(P)$ is μ -continuous because it is Scott-continuous and order-preserving. Hence $C\text{-Idl}(P)$ is a topological algebra with respect to the μ -topology as well, and hence the μ -closure $\bar{\bar{P}}$ of $\eta(P)$ is also a subalgebra. By Proposition 2.9 it contains \bar{P} .

4 Inequations

We are ready to embark on the main result of this paper, namely, that the free dcpo algebra \overline{P} satisfies all inequations that are valid in the preordered algebra P . The setting is slightly more general than in the previous section: we assume we are given a dcpo presentation $(P; \overline{\omega}, C)$ and a signature Ω of operation symbols, each of finite arity. We further assume that P is a preordered Ω -algebra, which means that for every $\omega \in \Omega$ with arity n a monotone map $\omega_P: P^n \rightarrow P$ has been specified. (Below we will no longer distinguish between the function symbol ω and the concrete operation ω_P .) We also assume that C is stable with respect to all operations ω_P , $\omega \in \Omega$.

Definition 4.1 *For Ω a set of function symbols, an inequation is a first-order formula of the form $(\forall \vec{x})t_1 \sqsubseteq t_2$, where t_1 and t_2 are Ω -terms whose variables all appear in the vector \vec{x} .*

The notion of algebra (model) for an inequational theory can be defined in any preorder-enriched category with products. For the purposes of this paper, we are interested in the categories **Pre** (of preorders) and **dcpo**.

Proposition 4.2 *Let $(P; \overline{\omega}, \Omega)$ be a preordered Ω algebra and C a set of covers stable for all $\omega \in \Omega$. Let \overline{P} be the dcpo algebra presented by $(P; \overline{\omega}, C)$ with Scott-continuous operations $\overline{\omega}$ ($\omega \in \Omega$) as defined in the previous section. Then any inequation that is valid in P also holds in \overline{P} .*

Proof. Let $(\forall \vec{x})t_1 \sqsubseteq t_2$ be an inequation that is valid in P . The terms t_1 and t_2 define n -ary monotone operations on P , hence extend to n -ary Scott-continuous operations $\overline{t}_1, \overline{t}_2$ on \overline{P} by Proposition 3.5. The set X of tuples \vec{a} in \overline{P}^n for which $\overline{t}_1(\vec{a}) \sqsubseteq \overline{t}_2(\vec{a})$ holds, is a sub-dcpo of \overline{P}^n . It includes $(\eta(P))^n$ because the inequation holds in P and η is a monotone homomorphism by Corollary 3.4. By Proposition 2.8, X is the whole of \overline{P}^n . ▮

Note that the analogous statement for the ambient algebra $C\text{-Idl}(P)$ is false in general. An example is easily given. Let P be trivially ordered and carry a binary operation that satisfies $x * x = x$; let the set C of covers be empty. The set of C -ideals consists of all subsets of P . Clearly, forming the product $A * A$ of a subset $A \subseteq P$ with more than one element may contain elements not belonging to A .

Definition 4.3 *For Ω a set of function symbols, an inequational theory is simply a set \mathcal{E} of inequations over Ω . A preordered algebra supporting all operations in Ω is called a (Ω, \mathcal{E}) -algebra if it satisfies the inequations in \mathcal{E} . We often abbreviate the pair (Ω, \mathcal{E}) to \mathbb{T} .*

Theorem 4.4 *Let $\mathbb{T} = (\Omega, \mathcal{E})$ be an inequational theory and P a preordered \mathbb{T} -algebra. Let C be a set of covers stable for all operations in Ω . Then $\eta: P \rightarrow \overline{P}$ makes \overline{P} freely generated as a dcpo- \mathbb{T} -algebra that respects the \mathbb{T} -algebra structure of P and preserves the C -covers.*

Proof. That \overline{P} is a dcpo- \mathbb{T} -algebra follows from the preceding proposition. Freeness was shown in Theorem 3.6. ▮

The phenomenon reported in this theorem was first noted by Abramsky and Vickers for a special algebraic theory; it is the essence of their “coverage theorem” in [AV93]. We believe that it is the “general unifying account” that is asked for in [VT04, page 301]. We will examine how the various coverage theorems in the literature follow from it in Section 6 below. In the remainder of the present section we indicate how Theorem 4.4 could have been established by other means.

The d -topology $C\text{-Idl}(P)$ is a topological algebra with respect to the d -topology. The subalgebra $\eta(P)$ satisfies the inequation of P as it is a monotone homomorphic image of P . Since the d -topology is Hausdorff, the topological closure preserves the inequalities. Finally, the closure is exactly \overline{P} .

The μ -topology Here we can not just invoke the topological closure argument because the μ -topology may not be Hausdorff. Still, the desired result holds:

Proposition 4.5 *The μ -closure $\overline{\overline{P}}$ of $\eta(P)$ satisfies the inequations of P .*

Proof. Let t_1, t_2 be two terms such that the corresponding term functions $\overline{t_1}, \overline{t_2}$ do not satisfy $\overline{t_1} \sqsubseteq \overline{t_2}$ on $\overline{\overline{P}}$. This means that there exists a vector \vec{I} of C -ideals, all taken from $\overline{\overline{P}}$, such that $\overline{t_1}(\vec{I}) \not\sqsubseteq \overline{t_2}(\vec{I})$. The operations are μ -continuous on $C\text{-Idl}(P)$ as they are Scott-continuous, and this extends to term functions. So we find approximating ideals \vec{I}' from $\eta(P)$ such that $\overline{t_1}(\vec{I}')$ belongs to the μ -open set $\downarrow \overline{t_1}(\vec{I}) \setminus \downarrow \overline{t_2}(\vec{I})$. It follows that $\eta(P)$ does not satisfy $(\forall \vec{x}) t_1 \sqsubseteq t_2$ either, and neither does P . \blacksquare

Iterated ideal completion Yet another way to establish our main theorem is via the transfinite generation process described at the end of Section 2. One first shows the following result directly (rather than relying on the fact that it is the special case $C = \emptyset$ in Proposition 4.2):

Proposition 4.6 *An inequational law that is valid in P also holds in $\text{Idl}(P)$.*

Then it is easy to see that inequations are preserved at every stage of the transfinite process.

5 Extensions

The phenomenon we are reporting in this paper is very robust and can be extended in a number of ways. First of all, the restriction to single-sorted algebras was purely for the sake of expository simplicity — the results of sections 3 and 4 hold equally well in the many-sorted case. Of course, one then deals with a dcpo presentation for each sort separately.

More interesting is the case where we are given an operation that takes values in $\text{Idl}(P)$ rather than P . This situation arises frequently in domain theory and also in the “flat site” coverage theorem at the end of this paper. We show how this case can be reduced to the standard one, so let $\omega: P^n \rightarrow \text{Idl}(P)$ be a monotone map. By setting

$$\omega'(\vec{A}) := \bigcup \{ \omega(\vec{a}) \mid \vec{a} \in \vec{A} \}$$

we obtain a monotone operation of the usual kind on the ordered set $\text{Idl}(P)$. The idea is now to complete $\text{Idl}(P)$, rather than P itself. We lift the given set C of covers to a set C' of covers on $\text{Idl}(P)$ through the following two rules:

$$\frac{(a \triangleleft U) \in C}{(\downarrow a \triangleleft \{\downarrow U\}) \in C'} \text{ (lift)} \qquad \frac{A = \bigsqcup_{i \in I} A_i \text{ in } \text{Idl}(P)}{(A \triangleleft \{A_i \mid i \in I\}) \in C'} \text{ (cont)}$$

Proposition 5.1 *The dcpo $\overline{\overline{P}}$ presented by $(P; \overline{\sqsubseteq}, C)$ is isomorphic to $\overline{\text{Idl}(\overline{P})}$ presented by $(\text{Idl}(P); \sqsubseteq, C')$.*

Proof. We show that $\overline{\text{Idl}(\overline{P})}$ has the required extension property. Consider the following diagram:

$$\begin{array}{ccccc} P & \xrightarrow{\quad} & \text{Idl}(P) & \xrightarrow{\eta} & \overline{\text{Idl}(\overline{P})} \\ & \searrow f & \downarrow f' & \swarrow \overline{f'} & \\ & & D & & \end{array}$$

where we assume that D is a dcpo and f preserves covers. The function f' is the unique Scott-continuous map for which $f = f' \circ \downarrow$. We show that it preserves the covers in C' . For covers created by the (lift)-rule, we compute:

$$\begin{aligned}
f'(\downarrow a) &= f(a) \\
(f \text{ preserves covers in } C) &\sqsubseteq \bigsqcup^\uparrow \{f(x) \mid x \in U\} \\
&= \bigsqcup^\uparrow \{f(x) \mid x \in \downarrow U\} \\
&= \bigsqcup^\uparrow \{f'(\downarrow x) \mid x \in \downarrow U\} \\
(f' \text{ is continuous}) &= f'(\bigsqcup^\uparrow \{\downarrow x \mid x \in \downarrow U\}) \\
&= f'(\downarrow U)
\end{aligned}$$

Covers created by the (cont)-rule are preserved because f' is continuous. It follows that the extension $\overline{f'}$ exists and satisfies

$$\overline{f'} \circ (\eta \circ \downarrow) = f' \circ \downarrow = f$$

If we had another continuous map $g: \overline{\mathbf{Idl}(P)} \rightarrow D$ with $g \circ (\eta \circ \downarrow) = f$, then $g \circ \eta$ would be another continuous map into D extending \downarrow and so would have to be equal to f' . (Here we are using that η is Scott-continuous, which is enforced by the covers created through the (cont)-rule.) However, $g \circ \eta = f'$ forces $g = \overline{f'}$ by the universal property of $\mathbf{Idl}(P)$. \blacksquare

In order to apply our theory of Section 3 we need to find a suitable requirement on C that ensures that C' is ω' -stable. Since ω' is Scott-continuous, stability for the covers created by the (cont)-rule is automatic.

Ensuring stability for the covers created by the (lift)-rule would lead to an awkward condition on ω , but luckily, the situation becomes much more malleable through the following concept:

Definition 5.2 For C a set of covers on a preorder $(P; \sqsubseteq)$ define the saturation $\mathbf{sat}(C)$ of C by the following rules:

$$\begin{aligned}
&\frac{(a \triangleleft U) \in C}{(a \triangleleft U) \in \mathbf{sat}(C)} \text{ (inc)} && \frac{a' \sqsubseteq a \quad (a \triangleleft U) \in \mathbf{sat}(C) \quad \downarrow U \subseteq \downarrow U'}{(a' \triangleleft U') \in \mathbf{sat}(C)} \text{ (mon)} \\
&\frac{(a \triangleleft U) \in \mathbf{sat}(C) \quad \forall b \in U. (b \triangleleft V) \in \mathbf{sat}(C)}{(a \triangleleft V) \in \mathbf{sat}(C)} \text{ (trans)}
\end{aligned}$$

It is clear that a monotone function that preserves the covers in C also preserves those in the saturation, and therefore that $(P; \sqsubseteq, C)$ and $(P; \sqsubseteq, \mathbf{sat}(C))$ present the same dcpo \overline{P} .

Definition 5.3 Let $\omega: P^n \rightarrow \mathbf{Idl}(P)$ be a monotone operation. A set of covers C is called ω -stable if whenever we have $p \in \omega(a_1, \dots, a_n)$ and $a_i \triangleleft U$ belongs to C , then $p \triangleleft U'$ for some $U' \subseteq \omega(a_1, \dots, U, \dots, a_n)$.

Proposition 5.4 If C is ω -stable for $\omega: P^n \rightarrow \mathbf{Idl}(P)$ then $\mathbf{sat}(C')$ is ω' -stable for $\omega': \mathbf{Idl}(P)^n \rightarrow \mathbf{Idl}(P)$.

Proof. Let $\downarrow a \triangleleft \{\downarrow U\}$ be a cover in C' created by the (lift)-rule. Given any vector of directed ideals $A_1, \dots, A_{i-1}, A_{i+1}, \dots, A_n$ we need to show that

$$\omega'(A_1, \dots, \downarrow a, \dots, A_n) \triangleleft \omega'(A_1, \dots, \downarrow U, \dots, A_n)$$

belongs to $\mathbf{sat}(C')$. This will follow from (cont) and (trans) if we can show

$$\downarrow p \triangleleft \omega'(A_1, \dots, \downarrow U, \dots, A_n)$$

for every $p \in \omega'(A_1, \dots, \downarrow a, \dots, A_n) = \bigcup_{a_k \in A_k} \omega(a_1, \dots, a, \dots, a_n)$. However, by ω -stability of C we have $p \triangleleft U'$ for some directed set $U' \subseteq \omega(a_1, \dots, U, \dots, a_n) \subseteq \omega'(A_1, \dots, \downarrow U, \dots, A_n)$ and the desired cover belongs to $\text{sat}(C')$ by rules (mon), (lift), and (cont). \blacksquare

To complete the translation from P to $\text{ldl}(P)$ observe that if an inequation $(\forall \vec{x})t_1 \sqsubseteq t_2$ is valid in P then it also holds in $\text{ldl}(P)$. By the extension Theorem 4.4 it will then also hold in the presented dcpo-algebra $\overline{\text{ldl}(P)}$.

6 Applications

6.1 Quotients and colimits in DCPO

Given a dcpo D we may wish to force certain inequalities to hold. The results of Section 2 show how this can be done, namely by collecting the inequations into a set of covers C , but they also show that the process of forming the quotient is non-trivial; this is because the naive (preorder) quotient may contain directed sets that did not exist before.

One application of quotienting is the construction of the coequalizer in **DCPO**. Coproducts are just disjoint unions, so together we have a fairly concrete description of colimits in **DCPO**.

6.2 Free dcpo algebras

Let \mathbb{T} be an inequational theory. One of our main results, Theorem 4.4, shows that from a \mathbb{T} -algebra $(P; \overline{\sqsubseteq})$ in **Pre**, equipped with a dcpo presentation that is stable for the operations in \mathbb{T} , the dcpo \overline{P} presented is a free \mathbb{T} -algebra in **dcpo** over P . In this section we shall use the result to construct, given a dcpo D , a free \mathbb{T} -algebra over D in **dcpo**. In other words, we are constructing a left adjoint to the forgetful functor $\text{Alg}_{\mathbb{T}}(\mathbf{dcpo}) \rightarrow \mathbf{dcpo}$. (For simplicity we are assuming here that \mathbb{T} is single-sorted. However, the extension to many-sorted theories is easy. It will give a left adjoint to the forgetful functor $\text{Alg}_{\mathbb{T}}(\mathbf{dcpo}) \rightarrow \mathbf{dcpo}^{\text{Sort}}$, where **Sort** is the set of sorts for \mathbb{T} .)

Let D be a dcpo. We shall assume D is presented as \overline{P} for some given dcpo presentation $(P; \overline{\sqsubseteq}, C)$. In a sense this is unnecessary, since D has a canonical presentation in which $P = D$ and C has a cover for every directed subset of D . However, there is some advantage in knowing how the construction can be made in terms of presentations. We proceed in a number of steps.

First, let $P_{\mathbb{T}}$ be the term algebra, the set of all terms made from elements of P and operators in \mathbb{T} .

Second, let $\overline{\sqsubseteq}_{\mathbb{T}}$ be the smallest congruence preorder on $P_{\mathbb{T}}$ that includes $\overline{\sqsubseteq}$ and satisfies the inequational laws in \mathbb{T} . By *congruence preorder*, we mean a preorder with respect to which the operations are monotone. That is to say, if $b \overline{\sqsubseteq}_{\mathbb{T}} c$ then $\omega(a_1, \dots, b, \dots, a_n) \overline{\sqsubseteq}_{\mathbb{T}} \omega(a_1, \dots, c, \dots, a_n)$ for any operator ω and elements a_i . $(P_{\mathbb{T}}; \overline{\sqsubseteq}_{\mathbb{T}})$ is a \mathbb{T} -algebra in **Pre**.

Third, let $C_{\mathbb{T}}$ be the smallest dcpo presentation on $P_{\mathbb{T}}$ that includes C and is stable for all the operators in \mathbb{T} .

Theorem 6.1 *Let $(P_{\mathbb{T}}, \overline{\sqsubseteq}_{\mathbb{T}}, C_{\mathbb{T}})$ be as above. Then $\overline{P}_{\mathbb{T}}$ is the free **dcpo** \mathbb{T} -algebra over D .*

Proof. Let E be a **dcpo** \mathbb{T} -algebra, and let $\overline{f} : D \rightarrow E$ be a continuous map. By definition of dcpo presentation, \overline{f} is equivalent to a monotone, cover-preserving function $f : P \rightarrow E$. This extends uniquely to a homomorphism $f_{\mathbb{T}} : P_{\mathbb{T}} \rightarrow E$. The inverse image of \sqsubseteq_E is a congruence preorder on $P_{\mathbb{T}}$ that includes $\overline{\sqsubseteq}$ and satisfies the inequational laws in \mathbb{T} , and it follows that $f_{\mathbb{T}}$ is monotone with respect to $\overline{\sqsubseteq}_{\mathbb{T}}$. Similarly, by considering those pairs $(a, U) \in P_{\mathbb{T}} \times \mathcal{P}P_{\mathbb{T}}$, with U directed, such that $f_{\mathbb{T}}(a) \sqsubseteq \bigsqcup_{u \in U} f_{\mathbb{T}}(u)$, we see that $f_{\mathbb{T}}$ preserves the $C_{\mathbb{T}}$ -covers. Hence by Theorem 4.4 $f_{\mathbb{T}}$ extends uniquely to a homomorphism $\overline{f}_{\mathbb{T}} : \overline{P}_{\mathbb{T}} \rightarrow E$. \blacksquare

6.3 Coverage theorems

The content of Theorem 4.4 is that it shows that the same structure (\overline{P}) can be presented by generators and relations in two different ways: either as a dcpo,

$$\mathbf{dcpo}\langle P \mid a \sqsubseteq \bigvee^\uparrow U (a \triangleleft U) \rangle$$

or as a dcpo- \mathbb{T} -algebra,

$$\mathbf{dcpo}\mathbb{T}\langle P \text{ (qua } \mathbb{T}\text{-algebra)} \mid a \sqsubseteq \bigvee^\uparrow U (a \triangleleft U) \rangle.$$

Presenting as a dcpo- \mathbb{T} would, other things being equal, create a larger structure since all the \mathbb{T} -terms have to be generated. But this is compensated for by the “qua \mathbb{T} -algebra”, in effect extra relations, which constrain the newly generated terms to relate to what already exists in P .

This is the typical pattern of the “coverage theorems” known from locale theory. In fact the underlying question is seen more widely in mathematics. One example from ring theory is the basic property of ideals. If R is a ring and I an ideal, then we have

$$\begin{aligned} \mathbf{Ring}\langle R \text{ (qua ring)} \mid a = 0 (a \in I) \rangle \\ \cong \mathbf{Ab}\langle R \text{ (qua Abelian group)} \mid a = 0 (a \in I) \rangle. \end{aligned}$$

The ideal property $RIR \subseteq I$ may be seen as a “multiplication stability” of the subgroup I .

The original locale-theoretic coverage theorem is that of [Joh82], which describes the construction of the frame of \mathbf{C} -ideals on a *site*, a meet-semilattice equipped with a meet-stable coverage (not necessarily a dcpo presentation). It was also shown that the frame of \mathbf{C} -ideals was freely generated over the semilattice, with respect to transforming covers to joins. In fact the constructions there underlie much of the present paper. However, something that was not explicit in [Joh82] was the fact that, even without the meets structure, the set of \mathbf{C} -ideals could be presented as a *sup-lattice* (complete join semilattice). This was stated and exploited in [AV93]. If (S, \mathbf{C}) is the site, then

$$\begin{aligned} \mathbf{Fr}\langle S \text{ (qua } \wedge\text{-semilattice)} \mid a \sqsubseteq \bigvee U (a \triangleleft U) \rangle \\ \cong \mathbf{SupLat}\langle S \text{ (qua poset)} \mid a \sqsubseteq \bigvee U (a \triangleleft U) \rangle, \end{aligned}$$

with both isomorphic to the set of \mathbf{C} -ideals. There are two directions in which this can be used. One is that if a frame is presented by generators and relations (in other words, it is the Lindenbaum algebra for a propositional geometric theory), then the presentation can be manipulated into site form and then the result shows how to define *sup-lattice* homomorphisms from the frame. This was particularly useful in the context of [AV93], which involved quantales and modules over them, which are sup-lattices but not frames in general. The other direction was that it shows how to generalize the coverage theorem to get presentations for structures other than frames, specifically quantales and their modules.


A specific aspect of the technique is that once the existence of the sup-lattice $\mathbf{SupLat}\langle S \text{ (qua poset)} \mid a \sqsubseteq \bigvee U (a \triangleleft U) \rangle$ is known, one can forget its concrete representation as a set of \mathbf{C} -ideals. Its universal property as sup-lattice can be used to define meet on it as a bilinear (with respect to joins) operation, and then show that the operation gives meet with respect to the sup-lattice order. This is similar to the way in which we here extend operations from the preorder to the dcpo.

In [JV91] those same ideas were developed with sup-lattices replaced by *preframes* – a preframe is a dcpo with finite meets, binary meet distributing over directed joins. It was shown how frame presentations in a certain form could be reduced to preframe presentations. This time,

the set of generators was a *join*-semilattice. The relations were all of the form $\bigwedge S \leq \bigvee_i^\uparrow \bigwedge T_i$, with a certain join-stability property. Again, once the presented preframe was known to exist (this was a substantial result of the paper), its frame structure could be proved from the universal properties.

[VT04] combines those “sup-lattice” and “preframe” coverage theorems to prove a “double” coverage theorem that relates frame presentations to dcpo presentations. In fact it is a direct corollary of our Theorem 4.4.

[VT04] summarizes the coverage theorems in a cubical diagram



Each arrow here represents a forgetful functor that has a left adjoint, a free algebra functor. Note that some of the arrows are between finitary theories, and there are already non-trivial coverage theorems for those. However, our main Theorem, 4.4, is about dcpos and the coverage theorems for the central square.

6.3.1 Immediate applications

The following two results in [VT04] are immediate corollaries of Theorem 4.4. Note that [VT04] deals with equations: each “directed relation” is an equation between two joins of directed families. Our inequational form is normally more general, but in the case of semilattices each inequation can be expressed as an equation.

Proposition 6.2 *Let P be a join semilattice and R a join-stable set of directed relations on it. Then*

$$\mathbf{Sup}\langle P \text{ (qua } \vee\text{-SemiLat)} \mid R \rangle \cong \mathbf{dcpo}\langle P \text{ (qua poset)} \mid R \rangle.$$

Proof. [VT04] says, “The standard technique applies.” What is meant by this is that once the RHS is known to exist then its universal property can be used to define \vee . The task then is to show that the operation so obtained turns the RHS into a sup-lattice, and to prove the sup-lattice universal property required by the left-hand side. Our Theorem 4.4 works differently; the operation \vee on P extends to an operation $\bar{\vee}$ for the general reasons explained in Section 3, and it satisfies all inequations that \vee satisfies. This is enough to show that $\bar{\vee}$ is the sup-operation on the ordered set \bar{P} , or more precisely, that $A \subseteq B$ holds if and only if $A\bar{\vee}B = B$ for all C -ideals $A, B \in \bar{P}$.

Assume $A \subseteq B$. Then $A\bar{\vee}B \subseteq B\bar{\vee}B = B$ by monotonicity and idempotence, and $B \subseteq A\bar{\vee}B$ holds because it holds for \vee on P .

Conversely, assume $A\bar{\vee}B = B$. Then $A \subseteq A\bar{\vee}B = B$, again using that $(\forall x, y)x \leq x \vee y$ holds in the sup-lattice P . ▮

Theorem 6.3 (Double Coverage Theorem) *If (P, R, \dots) is a DL-site, then*

$$\mathbf{Fr}\langle P \text{ (qua DL)} \mid R \rangle \cong \mathbf{dcpo}\langle P \text{ (qua poset)} \mid R \rangle$$

Proof. “DL-site” means that P is a distributive lattice and each relation in R is in the form of an equation between two joins of directed families in P . Moreover, the equation set is join- and meet-stable. Each equation can be expressed as a pair of inequations, so the whole site can be rephrased as a join- and meet-stable dcpo presentation.

The order on P is linked to the join operation as in the previous proposition, and to meet by equations. Hence the order on \bar{P} is linked to $\bar{\vee}$ as before and also to $\bar{\wedge}$ because the equations are preserved. \blacksquare

Proposition 6.4 *Let P be a meet semilattice and R a meet-stable set of directed relations on it. Then*

$$\mathbf{PreFr}\langle P \text{ (qua } \wedge\text{-SemiLat)} \mid R \rangle \cong \mathbf{dcpo}\langle P \text{ (qua poset)} \mid R \rangle.$$

Proof. The proof that the order on \bar{P} is given by $\bar{\wedge}$ is analogous to the one given for $\bar{\vee}$ above. \blacksquare

As a corollary of Proposition 6.2, we get the following. It illustrates a typical technique of expanding the generator set with operations that are preserved, and expanding the relation set to ensure stability.

Corollary 6.5 *Let P be a preorder and C a coverage on it (not necessarily directed). Then*

$$\mathbf{Sup}\langle P \text{ (qua preorder)} \mid a \sqsubseteq \bigvee U \text{ (} a \triangleleft U \text{)} \rangle$$

is isomorphic to

$$\mathbf{dcpo}\langle \mathcal{FP} \text{ (qua } \sqsubseteq_L\text{-preorder)} \mid B \cup \{a\} \sqsubseteq \bigvee^\uparrow \{B \cup U' \mid U' \in \mathcal{FU}\} \text{ (} B \in \mathcal{FP}, a \triangleleft U \text{)} \rangle.$$

Here \mathcal{FP} is the finite powerset, and the lower preorder \sqsubseteq_L is defined by $A \sqsubseteq_L B$ if for every $a \in A$ there is some $b \in B$ with $a \sqsubseteq b$.

Proof. One first shows that $\mathbf{Sup}\langle P \text{ (qua preorder)} \mid a \sqsubseteq \bigvee U \text{ (} a \triangleleft U \text{)} \rangle$ is isomorphic to

$$\mathbf{SupLat}\langle \mathcal{FP} \text{ (qua } \cup\text{-semilattice)} \mid B \cup \{a\} \sqsubseteq \bigvee^\uparrow \{B \cup U' \mid U' \in \mathcal{FU}\} \text{ (} B \in \mathcal{FP}, a \triangleleft U \text{)} \rangle.$$

This uses straightforward calculations with presentations. In one direction a maps to $\{a\}$, while in the other A maps to $\bigvee A$. The relations are \cup -stable, and then Proposition 6.2 can be applied. \blacksquare

6.3.2 The standard coverage theorems

The original coverage theorem was that if (S, C) is the site, then

$$\begin{aligned} \mathbf{Fr}\langle S \text{ (qua } \wedge\text{-semilattice)} \mid a \sqsubseteq \bigvee U \text{ (} a \triangleleft U \text{)} \rangle \\ \cong \mathbf{SupLat}\langle S \text{ (qua poset)} \mid a \sqsubseteq \bigvee U \text{ (} a \triangleleft U \text{)} \rangle. \end{aligned}$$

Using Corollary 6.5, the RHS can be reduced to a dcpo presentation in which the generators are \mathcal{FP} preordered by \sqsubseteq_L . These already form a distributive lattice, with $A \wedge B = \{a \wedge b \mid$

$a \in A, b \in B$ }, and this allows us to deduce by the same means that the dcpo presented can be given a frame presentation, which can then be proved isomorphic to the LHS. We do not give the details here, since the discussion in Section is similar but more general.

[JV91] proves a preframe coverage theorem.

Theorem 6.6 *Let S be a \vee -semilattice, and let R be a set of preframe relations of the form $\bigwedge A \sqsubseteq \bigvee_{i \in I}^\uparrow \bigwedge B_i$, where the B_i s form a directed family with respect to the upper order \sqsubseteq_U . ($A \sqsubseteq_U B$ if for every $b \in B$ there is some $a \in A$ with $a \sqsubseteq b$.) Suppose also that the set of relations is \vee -stable, in the sense that if $\bigwedge A \sqsubseteq \bigvee_{i \in I}^\uparrow \bigwedge B_i$ is in R , and $x \in S$, then the relation*

$$\bigwedge \{x \vee y \mid y \in A\} \sqsubseteq \bigvee_{i \in I}^\uparrow \bigwedge \{x \vee y \mid y \in B_i\}$$

is also in R . Then

$$\mathbf{Fr}\langle S \text{ (qua } \vee\text{-semilattice)} \mid R \rangle \cong \mathbf{PreFr}\langle S \text{ (qua poset)} \mid R \rangle.$$

Proof. We have $\mathbf{PreFr}\langle S \text{ (qua poset)} \mid R \rangle \cong \mathbf{PreFr}\langle \mathcal{FS} \text{ (qua } \cup = \wedge\text{-semilattice)} \mid R \rangle$, and $\mathcal{FS}/\sqsubseteq_U$ is a distributive lattice. The preframe presented is in turn isomorphic to

$$\begin{aligned} \mathbf{PreFr}\langle \mathcal{FS} \text{ (qua } \cup = \wedge\text{-semilattice)} \mid C \cup \bigcup_{j=1}^m A_j \sqsubseteq \bigvee_{j=1}^\uparrow \{C \cup \bigcup_{j=1}^m B_{i_j} \mid i_j \in I_j\} \\ (C \in \mathcal{FS}, \bigwedge A_j \sqsubseteq \bigvee_{i \in I_j}^\uparrow \bigwedge B_i \text{ in } R), \end{aligned}$$

and the relations here are both \wedge -stable and \vee -stable. We can now use Theorem 4.4 to see that

$$\begin{aligned} \mathbf{dcpo}\langle \mathcal{FS} \text{ (qua } \sqsubseteq_U\text{-preorder)} \mid C \cup \bigcup_{j=1}^m A_j \sqsubseteq \bigvee_{j=1}^\uparrow \{C \cup \bigcup_{j=1}^m B_{i_j} \mid i_j \in I_j\} \\ (C \in \mathcal{FS}, \bigwedge A_j \sqsubseteq \bigvee_{i \in I_j}^\uparrow \bigwedge B_i \text{ in } R) \end{aligned}$$

is isomorphic both to the preframe presented above and to the frame

$$\begin{aligned} \mathbf{Fr}\langle \mathcal{FS} \text{ (preserving } \cup \text{ and } \wedge) \mid C \cup \bigcup_{j=1}^m A_j \sqsubseteq \bigvee_{j=1}^\uparrow \{C \cup \bigcup_{j=1}^m B_{i_j} \mid i_j \in I_j\} \\ (C \in \mathcal{FS}, \bigwedge A_j \sqsubseteq \bigvee_{i \in I_j}^\uparrow \bigwedge B_i \text{ in } R) \end{aligned}$$

which in turn is isomorphic to $\mathbf{Fr}\langle S \text{ (qua } \vee\text{-semilattice)} \mid R \rangle$. ■

6.3.3 Flat sites

We discuss here the original sup-lattice coverage theorem in a slightly generalized form.

A *flat site* is defined in [Vic06] as a triple $(P, \sqsubseteq, \triangleleft_0)$ where (P, \sqsubseteq) is a preorder and \triangleleft_0 is a subset of $P \times \mathcal{P}P$ such that if $a \triangleleft_0 U$ and $b \sqsubseteq a$, then $b \triangleleft_0 V$ for some $V \subseteq b \downarrow U$. (We write $A \downarrow B$ for $(\downarrow A) \cap (\downarrow B)$, $\downarrow A$ for the down-closure of A .)

In fact, this is just re-notation for the inductively generated formal topologies of [CSSV03]. Each flat site presents a frame $\mathbf{Fr}\langle P, \sqsubseteq, \triangleleft_0 \rangle$, defined as

$$\begin{aligned} \mathbf{Fr}\langle P \text{ (qua preorder)} \mid 1 \sqsubseteq \bigvee P \\ a \wedge b \sqsubseteq \bigvee (a \downarrow b) \\ a \sqsubseteq \bigvee U \text{ (} a \triangleleft_0 U \text{)}. \end{aligned}$$

This generalizes the [Joh82] notion of site, though in a way that is already understood from the way Grothendieck topologies are used in topos theory. If P is a meet-semilattice then the condition on the coverage is equivalent to meet stability, and the first two relations given, together with “qua preorder”, are equivalent to preservation of finite meets of P .

Theorem 6.7

$$\mathbf{Fr}\langle P, \sqsupseteq, \triangleleft_0 \rangle \cong \mathbf{SupLat}\langle P \text{ (qua preorder)} \mid a \sqsubseteq \bigvee U \text{ (} a \triangleleft_0 U \text{)} \rangle.$$

Proof. By Corollary 6.5, the RHS in the statement is isomorphic to

$$\begin{aligned} \mathbf{dcpo}\langle \mathcal{FP} \text{ (qua } \sqsupseteq_L \text{-preorder)} \mid \\ B \cup \{a\} \sqsubseteq \bigvee^\uparrow \{B \cup U' \mid U' \in \mathcal{FU}\} \text{ (} B \in \mathcal{FP}, a \triangleleft_0 U \text{)} \rangle. \end{aligned}$$

\mathcal{FP} is already a semilattice, with \cup providing a join with respect to \sqsupseteq_L . But it also has a meet operation $\wedge : \mathcal{FP} \times \mathcal{FP} \rightarrow \mathbf{Idl}(\mathcal{FP})$,

$$\begin{aligned} A \wedge B &= \{S \in \mathcal{FP} \mid (\forall s \in S)(\exists a \in A)(\exists b \in B)s \in a \downarrow b\} \\ &= \{S \in \mathcal{FP} \mid S \sqsupseteq_L A, S \sqsupseteq_L B\}. \end{aligned}$$

Extending these to operations on $\mathbf{Idl}(\mathcal{FP})$, we find they make it a distributive lattice. Notably, $A \wedge (B_1 \vee B_2) = (A \wedge B_1) \vee (A \wedge B_2)$. For if $S \sqsupseteq_L A$ and $S \sqsupseteq_L B_1 \cup B_2$ then we can find $S = S_1 \cup S_2$ with $S_i \sqsupseteq_L B_i$, and this suffices to show $S \in (A \wedge B_1) \vee (A \wedge B_2)$.

The coverage as it stands is obviously \vee -stable. However, for \vee -stability we shall need to extend it. By induction on n one sees that the dcpo as presented is isomorphic to

$$\begin{aligned} \mathbf{dcpo}\langle \mathcal{FP} \text{ (qua } \sqsupseteq_L \text{-preorder)} \mid \\ B \cup \{a_1, \dots, a_n\} \sqsubseteq \bigvee^\uparrow \{B \cup U' \mid U' \in \mathcal{F}(\bigcup_{i=1}^n U_i)\} \\ \text{(} B \in \mathcal{FP}, n \geq 0, a_i \triangleleft_0 U_i \text{ for } 1 \leq i \leq n \text{)} \rangle. \end{aligned}$$

This is still \vee -stable. For \wedge -stability, suppose we have $B \cup \{a_1, \dots, a_n\} \triangleleft \{B \cup U' \mid U' \in \mathcal{F}(\bigcup_{i=1}^n U_i)\}$, deriving from $B \in \mathcal{FP}$ and $a_i \triangleleft_0 U_i$. Suppose $S \in C \wedge (B \cup \{a_1, \dots, a_n\}) = (C \wedge B) \vee (C \wedge \{a_1, \dots, a_n\})$, so $S = S_1 \cup S_2$ with $S_1 \sqsupseteq_L B$ and $S_2 \sqsupseteq_L \{a_1, \dots, a_n\}$. If $S_2 = \{b_1, \dots, b_m\}$ then for each j we have $b_j \sqsupseteq a_i$ for some i , so $b_j \triangleleft_0 V_j \subseteq b_j \downarrow U_i$. Then

$$\{B \cup V' \mid V' \in \mathcal{F}(\bigcup_{j=1}^m V_j)\} \subseteq C \wedge \{B \cup U' \mid U' \in \mathcal{F}(\bigcup_{i=1}^n U_i)\}$$

as required for \wedge -stability.

We can now apply Theorem 4.4 to see that this dcpo is isomorphic to

$$\begin{aligned} \mathbf{Fr}\langle \mathcal{FP} \text{ (qua } \sqsupseteq_L \text{-preorder)} \mid \vee, \wedge, 1 \text{ and } 0 \text{ preserved,} \\ B \cup \{a_1, \dots, a_n\} \sqsubseteq \bigvee^\uparrow \{B \cup U' \mid U' \in \mathcal{F}(\bigcup_{i=1}^n U_i)\} \\ \text{(} B \in \mathcal{FP}, n \geq 0, a_i \triangleleft_0 U_i \text{ for } 1 \leq i \leq n \text{)} \rangle, \end{aligned}$$

which in turn is isomorphic to $\mathbf{Fr}\langle P, \sqsupseteq, \triangleleft_0 \rangle$. ■

References

- [AJ94] S. Abramsky and A. Jung. Domain theory. In S. Abramsky, D. M. Gabbay, and T. S. E. Maibaum, editors, *Semantic Structures*, volume 3 of *Handbook of Logic in Computer Science*, pages 1–168. Clarendon Press, 1994.
- [AV93] S. Abramsky and S.J. Vickers. Quantales, observational logic and process semantics. *Mathematical Structures in Computer Science*, 3:161–227, 1993.
- [CSSV03] T. Coquand, G. Sambin, J. Smith, and S. Valentini. Inductively generated formal topologies. *Annals of Pure and Applied Logic*, 124:71–106, 2003.
- [HP79] M. C. B. Hennessy and G. D. Plotkin. Full abstraction for a simple parallel programming language. In J. Beçvar, editor, *Mathematical Foundations of Computer Science*, volume 74 of *Lecture Notes in Computer Science*. Springer Verlag, 1979.
- [Joh82] P.T. Johnstone. *Stone Spaces*. Number 3 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1982.
- [JV91] P.T. Johnstone and S.J. Vickers. Preframe presentations present. In A. Carboni, M.C. Pedicchio, and G. Rosolini, editors, *Category Theory – Proceedings, Como 1990*, number 1488 in *Lecture Notes in Mathematics*, pages 193–212. Springer-Verlag, 1991.
- [Mar00] Keye Martin. *A Foundation for Computation*. PhD thesis, Tulane University, July 2000.
- [PP04] G. D. Plotkin and J. Power. Computational effects and operations: an overview. In M. Escardó and A. Jung, editors, *Proceedings of the Workshop on Domains VI*, volume 73 of *Electronic Notes in Theoretical Computer Science*, pages 149–163. Elsevier Science Publishers B.V., 2004.
- [Vic06] Steven Vickers. Compactness in locales and in formal topology. *Annals of Pure and Applied Logic*, 137:413–438, 2006.
- [VT04] S.J. Vickers and C.F. Townsend. A universal characterization of the double powerlocale. *Theoretical Computer Science*, 316:297–321, 2004.
- [Wyl81] O. Wyler. Dedekind complete posets and Scott topologies. In B. Banaschewski and R.-E. Hoffmann, editors, *Continuous Lattices*, volume 871 of *Lecture Notes in Computer Science*, pages 384–389. Springer Verlag, 1981.